

SOKOTO STATE HEALTH INDICATORS- MICS 2021

S/N	INDICATOR	NATIONAL AVERAGE	SOKOTO STATE SCORE
	Early Childhood Mortality Rates		
1	Neonatal mortality rate (per 1,000 live birth)	34	34
2	Post-neonatal mortality rate (per 1,000 live birth)	28	70
3	Infant mortality rate (per 1,000 live birth)	63	104
4	Child mortality rate (per 1,000 live birth)	42	109
5	Under-five mortality rate (per 1,000 live birth)	102	202
	Adolescent birth rate and total fertility rate		
6	Adolescent birth rate: 15-19 years (number of births per 1,000 women)	75	104
7	Total fertility rate: women aged 15-49 years (number of births per 1,000 women)	4.6	5.2
	Early childbearing (young women)		
8	Percentage of women age 20-24 years who have had a live birth before age 18	21.1	33.2
	Trends in early childbearing (women)		
9	Percentage of women with a live birth before age 15	5.6	0.7
10	Percentage of women with a live birth before age 18	23.4	--
	Need and demand for family planning (currently married/in union)		
11	Unmet need for family planning (%)	23.9	12.1

	Antenatal care coverage		
12	No antenatal care (%)	21.1	58.8
	Neonatal tetanus protection		
13	Percentage of women who received at least 2 tetanus toxoid containing vaccine doses during the pregnancy of the most recent live birth	60.8	25.7
	Place of delivery		
14	Delivered in health facility (%)	49.0	14.2
	Assistance during delivery		
15	Delivery assisted by any skilled attendant (%)	50.7	14.5
	Infants weighed at birth		
16	Percentage of live births weighed at birth (%)	36.1	6.8
	Post-natal health checks for newborns		
17	No postnatal care visit (%)	76.0	71.2
	Location of first PNC (post-natal care) visit for newborns		
18	Location of first post-natal care visit for newborns – Home (%)	46.5	82.4
	Thermal care for newborns		
19	Percentage of children who were dried (wiped) after birth	80.1	51.5
20	Percentage of children who were given skin-to-skin contact with mother	6.1	2.8
	Post-natal health checks for mothers		
21	Health check following birth while in facility or at home (%)	58.2	28.3
22	No postnatal care visit (%)	85.3	79.8

	Location of first PNC visit for mothers		
23	Location of first PNC visit for mothers – Home (%)	59.1	79.4
	Post-natal health checks for mothers and newborns		
24	Percentage of post-natal health checks within 2 days of birth for newborns	62.4	46.0
25	Percentage of post-natal health checks within 2 days of birth for mothers	61.4	38.4
26	Percentage of post-natal health checks within 2 days of birth for both mothers and newborns	58.5	34.0
27	Percentage of post-natal health checks within 2 days of birth for neither mother nor newborn	34.8	49.5
	Vaccinations in the first years of life (%)		
28	BCG	74.1	33.7
29	Polio - At birth	70.2	28.7
30	Polio - OPV1	78.0	44.7
31	Polio - OPV2	68.4	31.2
32	Polio - OPV3	56.2	21.0
33	Polio – IPV	62.2	26.7
34	Polio - OPV3 & IPV	49.2	12.5
35	HepB at birth	67.2	22.5
36	DTP-HepB-Hib (1)	70.3	28.8
37	DTP-HepB-Hib (2)	64.6	18.4
38	DTP-HepB-Hib (3)	56.6	11.5
39	PCV (1)	69.1	21.8

40	PCV (2)	62.6	14.2
41	PCV (3)	54.7	10.0
42	Measles 1	60.3	19.1
43	Yellow Fever	58.9	17.1
44	Meningitis A (Conjugate)	54.9	19.5
45	Basic antigens	44.2	7.2
46	No vaccinations	17.8	50.7
	Vaccinations by background characteristics		
47	Percentage of children age 12-23 months with vaccination records seen	56.0	16.0
48	Percentage of children age 24-35 months who received Measles 2	37.5	18.5
49	Percentage of children age 24-35 months who received full vaccination – Basic antigens	44.0	10.0
50	Percentage of children age 24-35 months who received full vaccination - All antigens	20.6	4.8
51	Percentage of children age 24-35 months who received – No vaccinations	13.0	32.9
	Reasons vaccinations were not received		
52	Thought the child was fully immunized (%)	29.8	22.5
53	Unaware of need for immunization (%)	9.2	14.4
54	Unaware of need to return for 2nd or 3 rd dose (%)	8.0	10.0
55	Place and/or time of immunization unknown (%)	4.4	1.3

56	Postponed until another time (%)	6.7	4.4
57	Mother/ caretaker too busy (%)	10.9	6.3
58	Family problem, including illness of mother (%)	6.0	2.5
59	No faith in immunization (%)	9.8	23.8
60	Fear of side reactions (%)	10.8	20.6
61	Myths (%)	2.0	10.6
62	Believes there were contraindications (%)	2.5	6.3
63	Rumours (%)	0.9	5.6
	Reported disease episodes		
64	Percentage of children who in the last two weeks had an episode of diarrhoea	13.0	17.8
65	Percentage of children who in the last two weeks had symptoms of acute respiratory infections	2.0	1.0
66	Percentage of children who in the last two weeks had an episode of fever	24.6	26.2
	Care-seeking during diarrhoea		
67	No advice or treatment sought (%)	34.8	45.4
	Care-seeking for and antibiotic treatment of symptoms of acute respiratory infection		
68	No advice or treatment sought (%)	25.5	--
69	Percentage of children with symptoms of ARI in the last two weeks who were given antibiotics	49.9	--
	Care-seeking during fever		
70	No advice or treatment sought (%)	31.4	51.3
	Breastfeeding status		

71	Percent exclusively breastfed (Children aged 0-5 months)	34.4	30.7
72	Percent appropriately breastfed (Children age 0-23 Months)	53.9	55.8
	Infant and young child feeding (IYCF) practices		
73	Percent of children who received Minimum dietary diversity	31.1	29.9
74	Percent of children who received minimum meal frequency	34.4	26.2
75	Percent of children who received minimum acceptable diet	12.0	10.8